

Sexual dimorphism in *Carinatogecko heteropholis* (Minton, Anderson, and Anderson, 1970) (Sauria: Gekkonidae) from Ilam Province, western Iran

^{1,3}BEHZAD FATHINIA, ¹NASRULLAH RASTEGAR-POUYANI, AND ²HOSSEIN MOHAMADI

¹Department of Biology, Faculty of Science, Razi University, Kermanshah, IRAN ²Head of Natural History Museums and Genetic Resources, Department of Environmental Protection Organization, Tehran, IRAN

Abstract.—Sexual dimorphism is a widespread phenomenon in animals, but so far undocumented in *Carinatogecko heteropholis*. In this study, 52 specimens were collected in Karezan, Ilam province, western Iran. The uni- and multivariate analyses performed on the morphological data revealed that females are larger than males. All of the sexual differences were female-biased, except for the infralabial scales.

Key words. *Carinatogecko heteropholis*, sexual dimorphism, statistical analysis, morphology, Ilam Province, Iran

Citation: Fathinia, B., Rastegar-Pouyani, N., and Mohamadi, H. 2011. Sexual dimorphism in *Carinatogecko heteropholis* (Minton, Anderson, and Anderson, 1970) (Sauria: Gekkonidae) from Ilam Province, western Iran. *Amphib. Reptile Conserv.* 5(1):47-53(e27).

Introduction

The genus *Carinatogecko* Golubev and Szczerbak, 1981 comprises three species, the Iranian keel-scaled gecko, the Iraqi keel-scaled gecko, and Anderson's keel-scaled gecko: all of them are found on the Iranian Plateau (Szczerbak and Golubev 1996; Anderson 1999; Torki 2011). The Iraqi keel-scaled gecko, *Carinatogecko heteropholis* (Minton, Anderson, and Anderson 1970) is a small species; its type locality in Iran is western Zagros foothills (Anderson 1999; Fathinia 2007; Rastegar-Pouyani et al. 2007). It is hypothesized that the genus *Carinatogecko* has a double Iranian-Mesopotamian origin (Fathinia 2007).

Sexual dimorphism (SD) is a common and widespread phenomenon in the animal world (Andersson 1994). Sexual size dimorphism (SSD) explains the status in which the males and females differ in measured values of certain morphological characteristics. Sexual size dimorphism (SSD) has been extensively described in reptiles (Andersson 1994; Kuo 2009). Sexual dimorphism in animals is revealed in three different aspects: behavior, size, and shape (Selander 1972). Numerous surveys have been carried out on sexual dimorphism in lizards (Stamps 1983; Rocha 1996; Carothers 1984; Trivers 1976; Molina-Borja 2003; Baird et al. 2003; Verrastro 2004; Bruner et al. 2005; Kaliontzopoulou et al. 2007).

Differences in the selective forces acting on male versus female body size are the main causes of sex differences in adult body size of animals (Cox 2006). Sexual dimorphism in lizards may result from differences in

food resource partitioning and sexual differences in energy allocation to growth (Baird et al. 2003).

To our knowledge this is the first survey on the occurrence of sexual dimorphism in the genus *Carinatogecko*. Clarifying the sexually distinctive traits in *C. heteropholis* is of evolutionary and systematic importance; in this paper, we report results of such a study.

Material and methods

A total of 52 (28♂ and 24♀) adult specimens were collected during summer 2010. All of them were collected by hand with the aid of an electric torch at night on rocky mountain sides of the Zagros Mountains in Karezan, Shirvan-Chardavol, Ilam Province, western Iran (Fig. 1). Of these, 22 specimens were fixed in ethanol 75% and deposited in the RUZM (Razi University Zoological Museum) for future studies, and the rest (30 specimens) were released in their relevant habitat 24 hours after collecting and analyzing. The coordinates of the study site are 33°44' N, 46°29' E 1325 m a.s.l. Eight metric and four meristic variables were chosen and measured by digital caliper and stereomicroscope to the nearest 0.01 mm (Table 1). Except for overall shape differences which can be used to distinguish males from females (Fig. 2), sex of specimens was mainly determined based on presence of two swellings at the base of tail just behind vent in males and their absence in females (Fig. 3).

Correspondence. ³Email: bfathinia@gmail.com




Figure 1. Map showing the coordinates of the study site in Karezan region at mountainsides of the Zagros.

To determine the significance of sexual dimorphism in *C. heteropholis*, the ANOVA Table as well as Principal Component Analysis (PCA: correlation matrix) were used. The SPSS statistical software (version 13) was used for carrying out the statistical analyses.

Results

Twelve morphological characters (eight metric and four meristic) were included in the analysis. The values for

the metric and meristic characters as well as the direction of differences and the significant characters ($P < 0.05$) are summarized in Table 2.

ANOVA Table Analysis

Metric variables: obvious differences in the value of variables are observed between the sexes. Females have significantly greater values than the males for eight metric characters. In the case of body length and the distance




Figure 2. Dorsal view of male (left) and female (right) of *Carinatogecko heteropholis*.



Figure 3. Presence of swelling in the male of *C. heteropholis* at base of the tail which accommodate hemipenes (left) and their absence in female (right).




Figure 4. Ordination of the individual males and females of *Carinatogecko heteropholis* on the first two principal components. Note the relative degree of isolation between males and females, which is mainly attributed to SVL, TL, HL, HW, LFL, LHL, FHL, and VL in the PC1 and SL and IL in the PC2.

between forelimb – hindlimb (i.e., SVL and FHL, respectively) females had values of 36.34 ± 0.63 and 18.02 ± 0.41 and males had 32.53 ± 0.33 and 15.41 ± 0.20 ($P < 0.05$). Regarding the differences in extremities (forelimb, hindlimb, and tail) between females and males we observed that females had values of 12.96 ± 0.22 , 17.46 ± 0.30 , and 39.11 ± 0.80 and males had values of 11.86 ± 0.10 , 15.95 ± 0.19 , and 36.74 ± 0.68 for LFL, LHL, and TL respectively. Head dimensions also show significant differences between the sexes. Females had values of 8.94 ± 0.13 and 6.99 ± 0.10 and males had 8.51 ± 0.08 and 6.65 ± 0.07 for HL (head length) and HW (head width), respectively. Regarding the last metric character (i.e., VL or vent length), we realized that this character is significantly different between females and males, so that females have significantly greater values for VL ($3.58 \pm$

Table 1. The metric and meristic characters used in this study.

Characters	Definition
Metric	SVL
	snout to vent length
	TL
	length of tail
	HL
	head length
	HW
	head width
Meristic	LFL
	length of forelimb
	LHL
	length of hindlimb
FHL	FHL
	forelimb to hindlimb length
VL	VL
	the greatest horizontal length of vent
Meristic	SL
	number of supralabial scales
	IL
	number of infralabial scales
CT	CT
	number of crossbars on the tail
CD	CD
	number of chevrons on dorsum

0.10) than males (3.34 ± 0.04) ($P < 0.05$). All the metric variables are female biased. Reasons for presence of female biased sexual size dimorphism in the species are taken up in the discussion section.

Meristic variables: Significant differences were not observed in meristic variables, but SL (8.20 ± 0.12), CT (12.16 ± 0.24), and CD (7.45 ± 0.17) in females are larger than SL (8.07 ± 0.10), CT (11.96 ± 0.21), and CD (7.32 ± 0.14) in males. In other words, the three characters are not significantly female biased. Only one out of twelve variables (i.e., number of infralabials, IL) was male biased, which in turn was insignificant. The value of IL in males (6.85 ± 0.09) was insignificantly greater than that in females (6.79 ± 0.13) ($P < 0.05$).

Principal Component Analysis

The PCA performed on the dataset yielded three axes, which collectively explained 73.38% of the total variation. The PC1 explains 50.788% of the total variation. Inspection of the loadings indicates that correlations with all morphological measurements have the same sign (positive) but not the same magnitude (Table 3). The first axis is a clear indicator of body size. All metric variables in the first axis have greater values than meristic ones, hence making a greater contribution in sexual discrimination. The scores of the females along this axis show an overlap with those for males, indicating that although sexual dimorphism occurs between males and females, the two sexes are not completely separated from each other regarding these characters (Fig. 4). The second axis, which contains 12.51% of the total variation is a meristic axis that records individuals at one end with large SL and IL and relatively small SVL compared with individuals with small SL and IL and relatively large SVL. The third axis contains only 10.08% of the total variation, being a meristic axis that records individuals with large CT and CD and relatively small VL at one end, compared with individuals at the other end with small CT and CD and relatively high values for VL.

Discussion

Carinatogecko heteropholis presented marked sexual dimorphism in general body size and several body parts, with females being significantly larger than males in eight out of 12 studied characters.

The evolutionary result of selection acting differently on body size and the rest of male and female traits is sexual size dimorphism (SSD) (Andersson 1994). Both the proximate (growth patterns) and ultimate (evolutionary payoffs) causes are responsible for sexual dimorphism (Stamps 1993; Cox et al. 2003; Kuo et al. 2009). Regarding size dimorphism, the proximate cause is an agent which creates intersexual differences in growth rate. Among these proximate causes, two are mention-

Table 2. Comparison of 12 morphological characters in males and females of *Carinatogecko heteropholis*. SE: standard error, D of d: Direction of difference. All measurements in millimeter (mm). Abbreviations: SVL (snout-vent length), TL (length of tail), HL (head length), HW (head width), LFL (length of forelimb), LHL (length of hindlimb), FHL (forelimb-hindlimb length), VL (the greatest horizontal length of vent), SL (number of supralabial scales), IL (number of infralabial scales), CT (number of crossbars on the tail), and CD (number of chevrons on dorsum).

SEX	SVL	TL	HL	HW	LFL	LHL	FHL	VL	SL	IL	CT	CD
♂	Mean	32.53	36.74	8.51	6.65	11.86	15.95	15.41	3.34	8.07	6.85	11.96
	N	28	21	28	28	28	28	28	28	28	28	28
	SEM	0.33	0.68	0.08	0.07	0.10	0.19	0.20	0.04	0.10	0.09	0.21
♀	Mean	36.34	39.11	8.94	6.99	12.96	17.46	18.02	3.58	8.20	6.79	12.16
	N	24	24	24	24	24	24	24	24	24	24	24
	SEM	0.63	0.80	0.13	0.10	0.22	0.30	0.41	0.10	0.12	0.13	0.24
D. of d.		F>M	M>F	F>M	F>M							
P-value		0.000	0.027	0.008	0.009	0.000	0.000	0.000	0.030	0.386	0.691	0.530

able: differences in growth hormone concentrations and trade-offs in allocating energy between growth and reproduction (John-Adler et al. 2007; Kuo et al. 2009). Presence of dimorphism between males and females are defined by three main forces including: sexual, fecundity, and natural selection (Olsson et al. 2002; Cox et al. 2003; Kaliontzopoulou et al. 2007).

Ectotherms grow continuously throughout life and they show a tendency to produce abundant, varying numbers of progeny, which results in a vigorous correlation between fecundity and body size of females, and

probably that is why SSD in ectotherms is predominantly female-biased (Trivers 1972). The SVL (snout-vent length) and FHL (forelimb to hindlimb length) in females of *C. heteropholis* are greater than those in males. In other words, the two characters, SVL and FHL, are female-biased which can be the result of fecundity selection in the species. A larger abdominal volume is an ultimate cause which is selected in females because this feature enhances fecundity (Monnet and Cherry 2002, Tague 2005; Kuo et al. 2009).

Head size in a variety of lizards is male-biased (e.g. Verrastro 2004; Smith and Nickel 2002; Vial and Stewart 1989; Anderson and Vitt 1990; Castilla and Bauwens 1991; Mouton and van Wyk 1993; Vitt and Colli 1994; Barbadillo et al. 1995; Hews 1996; Smith et al. 1997; Shine et al. 1998; Kratochvíl and Frynta 2002). In the cases of HL (head length) and HW (head width) in *C. heteropholis*, females have significantly greater values than males. As reported for other vertebrates, a phenomenon which can support niche divergence hypothesis is dimorphism in head size (Selander 1972; Shine 1989). Reproductive role hypothesis is a hypothesis that explains differences in head size. Females have a greater contribution in reproduction (Darwin 1871) and a larger head should maximize energy intake. This idea may explain the presence of larger heads in females of *C. heteropholis*.

Further, in *C. heteropholis*, the volumes of LFL (length of forelimb) and LHL (length of hindlimb) in females are significantly greater than in males. Sexually size-adapted dimorphism in traits such as head, limb, and tail measurements are assigned to an artifact of the acceptance of SVL for scaling to body size (Kratochvíl et al. 2003). Moreover, we suggest that longer and stronger limbs are necessary to support greater distance between forelimb and hindlimb (i.e., greater FHL) either in females or in males.

Our results show that in the case of *C. heteropholis* the VL (vent length) in females is significantly greater than in males. During mating, females with a larger VL

Table 3. Loadings from a Principal Component Analysis of metric and meristic characters of *Carinatogecko heteropholis*. Variables loading strongly on each principal component are in bold. Abbreviations: SVL (snout-vent length), TL (length of tail), HL (head length), HW (head width), LFL (length of forelimb), LHL (length of hindlimb), FHL (forelimb-hindlimb length), VL (the greatest horizontal length of vent), SL (number of supralabial scales), IL (number of infralabial scales), CT (number of crossbars on the tail), and CD (number of chevrons on dorsum).

Variable	PC1	PC2	PC3
SVL	0.958	-0.133	-0.054
TL	0.791	0.061	0.003
HL	0.884	-0.057	-0.042
HW	0.848	-0.087	-0.032
LFL	0.907	-0.055	-0.031
LHL	0.911	-0.073	-0.028
FHL	0.833	-0.066	-0.007
VL	0.756	0.254	-0.195
SL	0.147	0.784	-0.064
IL	0.102	0.851	0.077
CT	0.286	-0.156	0.748
CD	0.140	0.174	0.771
Eigenvalue	6.095	1.502	1.210
% Variance	50.788	12.513	10.085
Cumulative	50.788	63.301	73.386

are chosen by males. According to Andersson (1994), this character in geckos may be the result of selection for fecundity as well as selection for a larger female VL during evolution.

Additional studies are needed to determine which of these alternatives best explain the occurrence of sexual dimorphism in *C. heteropholis*.

Acknowledgments.—We thank Hamzeh Oraei (Ph.D. student), Department of Biology, Faculty of Science, Tehran University, for his assistance in the statistical analysis and Rasoul Karamiani (M.Sc. student), Department of Biology, Faculty of Science, Razi University, for his contribution in identifying the species (i.e., *Carinatogecko heteropholis*).

References

- Anderson, R. A. and Vitt, L. J. 1990. Sexual selection versus alternative causes of sexual dimorphism in teiid lizards. *Oecologia* **84**(2):145-157.
- Anderson, S. C. 1999. *The Lizards of Iran*. Society for the Study of Amphibians and Reptiles, Ithaca, New York, USA. 442 p.
- Andersson, M. 1994. *Sexual Selection*. Princeton University Press, Princeton, New Jersey. 624 p.
- Baird, T. A., Vitt, L. J., Baird, T. D., Cooper, W. E., Jr., Caldwell, J. P., and Pérez-Mellado, V. 2003. Social behavior and sexual dimorphism in the Bonaire whiptail, *Cnemidophorus murinus* (Squamata: Teiidae): the role of sexual selection. *Canadian Journal of Zoology* **81**(11):1781-1790.
- Barbadillo, L. J., Bauwens, D., Barahona, F., and Sánchez-Herráiz, N. J. 1995. Sexual differences in caudal morphology and its relation to tail autotomy in lacertid lizards. *Journal of Zoology (London)* **236**(1):83-93.
- Bruner, E., Costantini, D., Fanfani, A., and Dell'Osso, G. 2005. Morphological variation and sexual dimorphism of the cephalic scales in *Lacerta bilineata*. *Acta Zoologica (Stockholm)* **86**(4):245-254.
- Carothers, J. H. 1984. Sexual selection and sexual dimorphism in some herbivorous lizards. *American Naturalist (Chicago)* **124**(2):244-254.
- Castilla, A. M. and Bauwens, D. 1991. Observations on the natural history, present status, and conservation of the insular lizard *Podarcis hispanica atrata* on the Colubretes Archipelago, Spain. *Biological Conservation* **58**(1):69-84.
- Cox, R. M. 2006. A test of the reproductive cost hypothesis for sexual size dimorphism in Yarrow's spiny lizard *Sceloporus jarrovii*. *Journal of Animal Ecology* **75**(6):1361-1369.
- Cox, R. M., Skelly, S. L., and John-Alder, H. B. 2003. A comparative test of adaptive hypotheses for sexual size dimorphism in lizards. *Evolution* **57**(7):1653-1669.
- Darwin, C. 1871. *The Descent of Man, and Selection in Relation to Sex*. John Murray, London. Volume 1: 423 p., Volume 2: 476 p.
- Fathinia, B. 2007. *The Biosystematic Study of Lizards of Ilam Province*. M.Sc. Thesis, Lorestan University. 126 p.
- Hews, D. K. 1996. Size and scaling of sexually-selected traits in the lizard, *Uta palmeri*. *Journal of Zoology (London)* **238**(4):743-757.
- Kaliotzopoulou, A., Carretero, M. A., and Llorente, G. A. 2007. Multivariate and geometric morphometrics in the analysis of sexual dimorphism variation in *Podarcis* lizards. *Journal of Morphology* **268**(2):152-165.
- Kratochvíl, L. and Frynta, D. 2002. Body size, male combat and the evolution of sexual dimorphism in eublepharid geckos (Squamata: Eublepharidae). *Biological Journal of the Linnaean Society* **76**(2):303-314.
- Kratochvíl, L., Fokt, M., Rehák, I., and Frynta, D. 2003. Misinterpretation of character scaling: a tale of sexual dimorphism in body shape of common lizards. *Canadian Journal of Zoology* **81**(6):1112-1117.
- Kuo, C., Lin, Y., and Lin, Y. 2009. Sexual size and shape dimorphism in an Agamid lizard, *Japalura swinhonis* (Squamata: Lacertilia: Agamidae). *Zoological Studies* **48**(3):351-361.
- Molina-Borja, M. 2003. Sexual dimorphism of *Gallotia atlantica atlantica* and *Gallotia atlantica mahoratae* (Lacertidae) from the eastern Canary Islands. *Journal of Herpetology* **37**(4):769-772.
- Monnet, J. M. and Cherry, M. I. 2002. Sexual size dimorphism in anurans. *Proceedings of the Royal Society of London Series B - Biological Sciences* **269**(1507):2301-2307.
- Mouton, P. le F. N. and van Wyk, J. H. 1993. Sexual dimorphism in cordylid lizards: a case study of the Drakensberg crag lizard, *Pseudocordylus melanotus*. *Canadian Journal of Zoology* **71**(9):1715-1723.
- Olsson, M., Shine, R., Wapstra, E., Ujvari, B., and Madsen, T. 2002. Sexual dimorphism in lizard body shape: the roles of sexual selection and fecundity selection. *Evolution* **56**(7):1538-1542.
- Rastegar-Pouyani, N., Johari, S. M., and Rastegar-Pouyani, E. 2007. *Field Guide to the Reptiles of Iran. Volume 1: Lizards*. 2nd edition. Kermanshah. 119 plates + 139 p. (In Persian).
- Rocha, C. F. D. 1996. Sexual dimorphism in the sand lizard *Liolaemus lutzae* of southeastern Brazil, p.131-140 in Péfaur, J. E. (editor). *Herpetología Neotropical. Actas del II Congreso Latinoamericano de Herpetología Mérida*, Universidad de los Andes, Conselho de Publicaciones, Volume 2.
- Selander, R. K. 1972. Sexual selection and dimorphism in birds, p. 180-230 in Campbell, B. (Editor). *Sexual Selection and the Descent of Man, the Darwinian*

- Pivot. Heinemann Educational Books, London. 388 p.
- Shine, R. 1989. Ecological causes for the evolution of sexual dimorphism: a review of the evidence. *The Quarterly Review of Biology* **64**(4):419-431.
- Shine, R., Keogh, S., Doughty, P., and Giragossyan, H. 1998. Costs of reproduction and the evolution of sexual dimorphism in a ‘flying lizard’ *Draco melanopogon* (Agamidae). *Journal of Zoology* (London) **246**(2):203-213.
- Smith, G. R., Lemos-Espinal, J. A., and Ballinger, R. E. 1997. Sexual dimorphism in two species of knob-scaled lizards (genus *Xenosaurus*) from Mexico. *Herpetologica* **53**:200-205.
- Smith, G. R. and Nickle, A. M. 2002. Sexual dimorphism in three Cuban species of curly-tailed lizards (*Leiocephalus*). *Caribbean Journal of Science* **38**(1-2):140-142.
- Stamps, J. A. 1983. Sexual selection, sexual dimorphism and territoriality in lizards, p.169-204 in Huey, R. B., Pianka, E. R., and Schoener, T. W. (editors). *Lizard Ecology: studies on a model organism*. Harvard University, Cambridge, Massachusetts. 512 p.
- Stamps, J. A. 1993. Sexual size dimorphism in species with asymptotic growth after maturity. *Biological Journal of the Linnean Society* **50**(2):123-145.
- Tague, R. G. 2005. Big-bodied males help us recognize that females have big pelvis. *American Journal of Physical Anthropology* **127**(4):392-405.
- Torki, F. 2011. Description of a new species of *Carinatogecko* (Squamata: Gekkonidae) from Iran. *Salamandra* **47**(2):103-111.
- Trivers, R. L. 1972. Parental investment and sexual selection, p. 136-179 in Campbell, B. G. (editor). *Sexual Selection and the Descent of Man, the Darwinian Pivot*. Heinemann Educational Books, London. 388 p.
- Trivers, R. L. 1976. Sexual selection and resource-accruing abilities in *Anolis garmani*. *Evolution* **30**:253-269.
- Verrastro, L. 2004. Sexual dimorphism in *Liolemus occipitalis* (Iguania, Tropiduridae). *Iheringia. Série Zoologia* **94**(1):45-48. [Online]. Available: http://www.scielo.br/scielo.php?pid=S0073-47212004000100007&script=sci_arttext [Accessed: 14 October 2011].
- Vial, L. J. and Stewart, J. R. 1989. The manifestation and significance of sexual dimorphism in anguid lizards: a case study of *Barisia monticola*. *Canadian Journal of Zoology* **67**(1):68-72.
- Vitt, L. J. and Colli, G. R. 1994. Geographical ecology of a Neotropical lizard: *Ameiva ameiva* (Teiidae) in Brazil. *Canadian Journal of Zoology* **72**(11):1986-2008.



Behzad Fathinia earned his B.A. and M.S. from Isfahan and Lorestan universities, respectively. His M.S. research focused on “The Biosystematic Study of Lizards of Ilam Province.” At present, he is a Ph.D. student at Razi University, Kermanshah, western Iran under supervision of Nasrullah Rastegar-Pouyani, Mozafar Sharifi, and Eskandar Rastegar-Pouyani. His dissertation research involves ecology, phylogeography, molecular systematics, and population genetics of the Iranian viper *Pseudocerastes urarachnoides* in western Iran. He is also interested in other reptiles, especially snakes.



Nasrullah Rastegar-Pouyani earned his B.S. in Zoology from Razi University Kermanshah, Iran in 1986 and his M.S. in Zoology from Tehran University, Tehran, Iran in 1991, where he studied herpetology with the agamids as the central object. He started his Ph.D. in Gothenburg University, Sweden in 1994 under the advisement of Professor Göran Nilson and graduated in 1999, working on taxonomy and biogeography of Iranian Plateau agamids with *Trapelus* as the main object. His research interests include taxonomy and biogeography of the Iranian Plateau, the Middle East and Central Asian herpetofauna.



Hussein Mohamadi is the Head of Natural History Museums and genetic resources, Environmental Protection Organization, Tehran, Iran. He earned his B.A. and M.S. degrees from the Natural Resource College of Tehran University. His M.S. thesis was “The Ecological study of the marsh Crocodile, *Crocodylus palustris*, in Baluchistan.” He is now continuing his study as a Ph.D. student in environmental science, branch of science and research at Islamic Azad University. The subject of his Ph.D. thesis is “The assessment of changing trends and modeling of habitat preference in yellow Persian deer, *Dama dama mesopotamica*.”

Manuscript received: 21 January 2011

Accepted: 05 September 2011

Published: 05 November 2011